

TREATING YOUR INFECTION

Patient name

Your doctor or nurse recommends
that you self-care

Back-up antibiotic
prescription issued

Your infection	Usually lasts	How to treat yourself better for these infections, now and next time	When you should contact your GP practice or contact NHS 111
<input type="checkbox"/> Middle-ear infection	4 days	<ul style="list-style-type: none"> Have plenty of rest. Drink enough fluids to avoid feeling thirsty. Ask your local pharmacist to recommend medicines to help your symptoms and pain. Fever is a sign the body is fighting the infection and usually gets better by itself in most cases. You can use paracetamol (or ibuprofen) if you or your child are uncomfortable as a result of a fever. Other things you can do suggested by your GP or nurse: <p>.....</p> <p>.....</p> <p>.....</p> <p>.....</p>	<p>1. to 8. are possible signs of serious illness and should be assessed urgently. Phone for advice if you are not sure how urgent the symptoms are.</p> <ol style="list-style-type: none"> If you develop a severe headache and are sick. If your skin is very cold or has a strange colour, or you develop an unusual rash. If you feel confused or have slurred speech or are very drowsy. If you have difficulty breathing. Signs can include: <ul style="list-style-type: none"> breathing quickly turning blue around the lips and the skin below the mouth skin between or above the ribs getting sucked or pulled in with every breath. If you develop chest pain. If you have difficulty swallowing or are drooling. If you cough up blood. If you are feeling a lot worse. <p>Less serious signs that can usually wait until the next available GP appointment:</p> <ol style="list-style-type: none"> If you are not improving by the time given in the 'Usually lasts' column. In children with middle-ear infection: if fluid is coming out of their ears or if they have new deafness. Other
<input type="checkbox"/> Sore throat	7 days		
<input type="checkbox"/> Common cold	10 days		
<input type="checkbox"/> Sinusitis	18 days		
<input type="checkbox"/> Cough or bronchitis	21 days		
<input type="checkbox"/> Other infection: days		

Back-up antibiotic prescription to be collected after days only if you do not feel better or you feel worse.

Collect from: GP reception GP or nurse Pharmacy

- Colds, most coughs, sinusitis, ear infections, sore throats, and other infections often get better without antibiotics, as your body can usually fight these infections on its own.
- The more we use antibiotics, the greater the chance that bacteria will become resistant to them so that they no longer work on our infections.
- Antibiotics can cause side effects such as rashes, thrush, stomach pains, diarrhoea, reactions to sunlight, other symptoms, or being sick if you drink alcohol with metronidazole.
- Find out more about how you can make better use of antibiotics and help keep this vital treatment effective by visiting [nhs.uk/antibiotics2017](https://www.nhs.uk/antibiotics2017)

Never share antibiotics and always return any unused antibiotics to a pharmacy for safe disposal.
Leaflet developed in collaboration with expert medical bodies.

Keep Working