

- **How do I store my tablets?**

Do not take your tablets if the expiry date has passed. Keep the tablets in a dry place below 25°C. Keep all medicines well out of the reach of children – it is best to lock them in a cupboard or medicine cabinet. Edoxaban may be re-dispensed in a compliance aid or blister pack.

- **Will edoxaban interact with other medicines, food or alcohol?**

Edoxaban has fewer potential interactions with other medicines compared with warfarin, and at present there are no known interactions with specific foods or alcohol. There are some medicines that edoxaban does interact with so patients should inform their prescriber of the names of all medicines they are taking (including prescription and over-the-counter medicines, vitamins and herbal supplements).

- **Should patients stop taking edoxaban if they are going to have a dental or medical procedure?**

Patients should not stop taking edoxaban without first talking to their doctor or dentist. Edoxaban may need to be stopped for one or more days before any planned surgery, dental or medical procedure.

All patients should be given the edoxaban alert card and counselled on the details. This should be carried with them at all times.

Date of preparation November 2015, reviewed June 2018, next review June 2021.

Acknowledgment and thanks: Countess of Chester Hospital

PATIENT INFORMATION LEAFLET

EDOxabAN FOR THE TREATMENT OF NON-VALVULAR ATRIAL FIBRILLATION

People with atrial fibrillation (AF) are at increased risk of having a stroke. If you have AF, your doctor should assess your individual risk of a stroke and discuss with you as to whether you should be started on medication to reduce this risk. This leaflet has been devised to provide information in conjunction with the advice from your doctor.

- **What is edoxaban for?**

Edoxaban (Lixiana®) is an anticoagulant (like warfarin) used to lower the risk of blood clots developing in patients who have a condition called atrial fibrillation (an abnormal heartbeat) as well as other risk factors for stroke. A blood clot which blocks an artery (blood vessel) is called an embolism. If the embolism occurs in the arteries of the brain it can cause a stroke. Anticoagulants slow down the time it takes for your blood to clot and reduce the risk of these events happening.

- **For patients with non-valvular atrial fibrillation, is it worth changing from warfarin?**

Warfarin has been prescribed for more than 60 years so there is plenty of experience of its clinical use. The clinical trial showed that edoxaban is as effective as warfarin. For patients whose warfarin is well controlled, it is probably not advisable to change. For patients who have poor anticoagulant control, a switch to edoxaban might be considered.

- **Does edoxaban cause less bleeding than warfarin?**

As both edoxaban and warfarin affect blood clotting, patients may still experience side effects such as bruising and bleeding. Intracranial bleeding (bleeding into the brain) is worrying because it is usually very serious. In the clinical trial, edoxaban caused less intracranial bleeding than warfarin.

Gastrointestinal (stomach and bowel) bleeding is also a concern as it varies widely in terms of severity and is more common. Edoxaban had significantly higher rates of gastrointestinal bleeding than warfarin.

- **If a patient has excessive bleeding, can the anticoagulant effect of edoxaban be reversed?**

There is no licensed product currently available to reverse bleeding with edoxaban. However, if urgent treatment is required, edoxaban will be discontinued and supportive measures will be started. It is easier to manage major bleeding in patients on warfarin.

- **Are regular blood tests needed to monitor edoxaban levels?**

There is no need for regular blood tests to measure the level of anticoagulant control with edoxaban. However, a blood test is needed to measure how well the kidneys are working before starting treatment and then at least once a year whilst on edoxaban treatment.

- **When and how do I take this medicine?**

Always take this medicine as your doctor has told you. If you do not take edoxaban regularly it will not be effective and your risk of a clot forming will increase.

The dose is usually 60mg once a day. This dose may be reduced to 30mg once a day if you have kidney problems, are taking other medicines which interact or have a higher risk of bleeding. Your doctor will discuss this with you.

Edoxaban can be taken with or without a meal preferably with water. Edoxaban can be re-dispensed into a compliance aid or blister pack.

- **What are the possible side effects of this medicine?**

As this medicine affects blood clotting, most side effects are related to signs such as bruising or bleeding. Tell your doctor immediately if you experience any of the following side effects: Long or excessive bleeding, exceptional weakness, tiredness, paleness, dizziness, headache or unexplained swelling. Also report if there is any blood in the urine or passing black tarry or blood-stained stools.

Patients should seek urgent medical attention if they fall or injure themselves during treatment, especially if they hit their head, due to the increased risk of bleeding.

- **What should I do if I forget to take a dose?**

It is important not to skip doses. If a dose is missed the patient should take edoxaban immediately and continue on the following day with the once daily intake as recommended. The dose should not be doubled within the same day to make up for a missed dose.

- **What should I do if I take too many tablets?**

If you have taken too many tablets all at once, you may have an increased risk of bleeding. Contact your doctor or local hospital casualty department at once. Show the doctor your pack of tablets.